

STELLAR COLLISIONS AND BLUE STRAGGLER STARS IN DENSE GLOBULAR CLUSTERS

SOURAV CHATTERJEE

Department of Astronomy, University of Florida, Gainesville, FL 32611.

FREDERIC A. RASIO

Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA) and
Dept. of Physics & Astronomy, Northwestern University, 2145 Sheridan Rd, Evanston, IL 60208, USA.

ALISON SILLS

Department of Physics and Astronomy, McMaster University, 1280 Main Street West, Hamilton, ON, L8S 4M1, CANADA.

AND

EVERT GLEBBEEK

Department of Astrophysics/IMAPP, Radboud University Nijmegen, PO Box 9010, 6500 GL, Nijmegen, The Netherlands.

Submitted to ApJ March 1, 2013

ABSTRACT

Blue straggler stars (BSS) are abundantly observed in all Galactic globular clusters (GGC) where data exist. However, observations alone cannot reveal the relative importance of various formation channels or the typical formation times for this well studied population of non-standard stars. Using a state of the art Hénon-type Monte Carlo code that includes all relevant physical processes, and observationally motivated initial conditions in large ranges in density, concentration, binary fraction, and mass, we create a little over 100 models with properties typical of the observed GGCs. These models can be directly compared with the observed GGCs and BSSs can be easily identified. We identify the BSSs in our models and determine their formation channels and times of creation. We find that for central densities above $10^3 M_{\odot} \text{pc}^{-3}$ the dominant formation channel is stellar collisions. The majority of these collisions happen via binary-mediated strong scattering encounters. As a result a strong correlation between the total number of BSSs and the binary fraction in a cluster can be observed. We find that the number of BSSs in the core show weak correlation with the collision rate Γ . Close inspection of our models reveals that the observed lack of correlation between the number of BSSs and Γ for the observed GGCs may be because the estimate of Γ using observed cluster properties is not a reliable indicator of the true Γ . Our models indicate that the BSSs observed today may have formed many Gyrs ago. Denser clusters have higher contributions from collisions which typically create higher mass BSSs compared to those created via mass-transfer in a binary. As a result, we find that the BSSs in denser clusters typically have formed more recently than those in sparser clusters.

Subject headings: methods: numerical — methods: statistical — blue stragglers — stars: kinematics and dynamics — globular clusters: general

1. INTRODUCTION

Blue straggler stars (BSSs) are stars bluer and brighter (and therefore more massive) than the main-sequence turnoff (in a cluster; Sandage 1953). It is impossible for normal single star evolution to create them. Instead, blue stragglers must be created when a main-sequence (MS) star is disturbed either via mass transfer in a binary or by a collision with another star. Both processes can increase the stellar mass and possibly also rejuvenate the star by bringing in new Hydrogen for burning in the core (eg., Lombardi et al. 1995, 1996; Sills et al. 1997, 2001; Lombardi et al. 2002). Since the BSSs are among the brighter members of a cluster, they are relatively easier to detect in photometric observations. All Galactic globular clusters (GGCs) where the data exist have shown this exotic stellar population in abundance (e.g., Piotto et al. 2002; Ferraro et al. 1995). Many open clusters also show moderately large populations (~ 10 s) of BSSs (e.g., Mathieu & Geller 2009).

A large body of theoretical work has been devoted to explaining the formation and properties of BSSs. This work suggest that there are two different formation channels for

the BSSs, namely collisions involving at least one MS star (Lombardi et al. 1995) and mass transfer in a binary (MTB) from a more evolved star (giant or another MS star) to a MS star (for a detailed recent study Chen & Han 2009). In both cases the key physical effect is replenishing the MS star with fresh Hydrogen (via a collision or via MTB), and increasing the star's mass. If this Hydrogen can be mixed inside the MS star and be brought to the core of the MS star for burning, then the MS star lives in the Hydrogen burning stage longer than an undisturbed star of equal stellar mass is expected to live. The star is then called rejuvenated. This rejuvenated star then would be observed as a BSS when its mass is larger than the standard turnoff mass in the cluster.

Stellar collisions can happen in various ways in a cluster. Two single stars can physically collide (SS collision). If binaries are present in the cluster, then binaries interact with other single stars (BS) or binaries (BB). During the BS and BB interactions collisions can occur between two or more stars taking part in the interactions (Fregeau et al. 2004).

In mass transferring binaries the details of the orbital evolution and mass transfer rate determines its final outcome. Mass transfer can be stable, where, a fraction of the companion

mass is transferred to the MS star. In contrast, dynamical mass transfer leads to merger of the two stars (Chen & Han 2009). Stellar mergers may also happen in interactions involving higher order hierarchical systems. For example, a hierarchical *triple* under certain configurations can increase the eccentricity of the inner binary orbit via Kozai effect (Kozai 1962). If the eccentricity is sufficiently high, the inner binary stars may be merged through tidal friction (Perets & Fabrycky 2009). Although it may be a significant channel of BSS formation in less dense open clusters (Perets & Fabrycky 2009), it is not clear whether these triples can be stable in more dense regions typical of the GGCs and thus be as effective.

Since BSSs are MS stars that have been disturbed, and possibly rejuvenated (by some channel), some form of interaction must have created them. These interactions are in turn dependent on the cluster global properties, including the central density (ρ_c), binary fraction (f_b), and mass-segregation timescale. Thus the BSSs in a cluster can in principle provide a window into the cluster’s dynamical history. In this respect the formation channel(s) of the BSSs and their typical formation times are of great interest. Unfortunately, although observations in various wavelength bands have taught us a lot about the stellar properties of the BSSs, it is hard to determine how a particular BSS formed or how long ago the interaction occurred. Theoretical modeling is the only way to learn this.

A popular approach so far to learn about the observed BSSs in the GGCs has been by carefully studying the correlations between the number of BSSs (normalized in some way by another characteristic stellar population in these clusters) with some dynamically important cluster property, including the collision rate estimator Γ (Pooley & Hut 2006), the binary fraction f_b (Sollima et al. 2008), and total cluster mass (most recently Leigh et al. 2013). While these observed correlations can teach us a lot about the possible formation channels of these exotic stellar populations, direct understanding of these processes are impossible to achieve. For example, a BSS created via mass transfer in a binary could actually be a result of some past dynamical encounter that had changed the orbital properties of a binary to alter its normal evolutionary path in such a way that mass transfer occurs. On the other hand, a collision in fact may be mediated by resonant dynamical interactions involving stellar binaries. In addition to these complicated interactions, it is also hard to know observationally exactly what the binary fraction in the cluster is now and is impossible to know what it had been in the past. Similarly, the calculation of the Γ parameter involves many properties in the cluster that are hard to determine. The only way to explore and probe these complicated interactions in an evolving cluster is by actual N -body modeling of a realistic cluster with realistic N , stellar mass function, and binary fraction and including all relevant physical processes such as two-body relaxation, strong encounters including physical collisions, binary-binary (BB) and binary-single (BS) interactions, stellar evolution, and Galactic tidal stripping. In these models if all encounters are documented in addition to the overall cluster properties and stellar properties at all times, then all necessary information can be obtained. BSSs can then be identified from a snapshot at any given time from these models and for them the full history from $t = 0$ can simply be looked up.

Until recently realistic modeling of GGCs was hard to do in a star-by-star detail including all relevant physical processes. Hence, idealized simulations have been used to study the formation channels and their radial distribution in the clus-

ter (Mapelli et al. 2006). However, Northwestern group’s recently modified Hénon-type Monte Carlo code CMC is well suited to quickly model GGCs including all relevant physical processes. Using this code it is now possible to model many clusters with star-by-star detail using a large range of observationally motivated initial parameters. For example, a recent study with CMC has explored a large range in parameter space and successfully created a population of ~ 200 realistic cluster models that have properties very similar to the observed GGCs (Chatterjee et al. 2010, 2013). These models include all relevant physical processes, and since they also have realistic N , these models can be directly compared with the observed GGCs without any need for scaling. In addition to the dynamical properties, single and binary stellar evolution properties are also followed in these models. Thus these simulations provide an unprecedented opportunity to understand the dominant dynamical processes responsible for BSS production in typical GGCs. In addition, since these models also span a large range in dynamically important global properties typically observed in the GGCs, these models also can shed light on the widely used correlations between the number of BSSs in a cluster and the cluster’s global properties and show what these correlations truly mean.

This study is the first in a series where these models are explored, BSSs are extracted, their dynamical history is obtained and two very basic questions are answered. What is the dominant physical process that creates the BSSs observed in the GGCs? What is the typical age since formation for the BSSs?

This study is organized as follows. In Section 2 we describe briefly our simulation setup, describe how the BSSs are identified in the models, and definitions for various categories of BSSs based on their formation channels. In Section 3 we show examples of the radial distribution for our model BSSs. We present the relative importance of the various formation channels in Section 4. The time elapsed since formation for the BSSs in our models is explored in Section 5. In Section 6 we investigate correlations between the specific frequency of the BSSs in the core with f_b and Γ . We summarize and conclude in Section 7.

2. NUMERICAL METHOD

We use our Hénon-type Cluster Monte Carlo code (CMC) to numerically model star clusters typical of the observed GGCs including realistic number (N) of stars and primordial binary fraction (f_b). We include all relevant physical processes namely two-body relaxation, strong interactions such as physical collisions and binary mediated scattering interactions, Galactic tidal stripping, and single and binary stellar evolution. This code is developed and tested rigorously over the past decade (Joshi et al. 2000, 2001; Fregeau et al. 2003; Fregeau & Rasio 2007; Chatterjee et al. 2010; Umbreit et al. 2012; Pattabiraman et al. 2013; Chatterjee et al. 2013). Using a large grid of initial conditions over the range of values typical of the observed young massive clusters we create 121 detailed star-by-star models and evolve them for ~ 12 Gyr. Our choice of initial conditions is based on observations of super star clusters. All our simulated clusters have initial virial radii between $r_v = 3 - 4$ pc. Initial N is varied between $4 - 10 \times 10^5$ stars. The positions and velocities of the stars (and center of mass for the binaries) are assigned according to King profiles with the dimensionless concentration parameter W_0 between 4–8. We vary the initial binary fraction, f_b , between 0.05–0.3. The stellar masses of the stars, or pri-

aries in case of a binary, are chosen from the IMF presented in Kroupa (2001, their Equations 1 and 2) in the stellar mass range $0.1 - 100 M_{\odot}$. Secondary binary companion masses are sampled from a uniform distribution of mass ratios in the range $0.1 M_{\odot} - m_p$, where m_p is the mass of the primary. The semi-major axes, a , for stellar binaries are chosen from a distribution flat in $\log a$ within physical limits, namely between $5 \times$ the physical contact of the components and the local hard-soft boundary. Although initially all binaries in our models are hard at their respective positions, some of these hard binaries can become soft during the evolution of the cluster. The cluster contracts under two-body relaxation and the velocity dispersion increases making initially hard binaries soft. Moreover, binaries sink to the core due to mass segregation where the velocity dispersion is higher than that at the initial binary positions. We include these soft binaries in our simulations until they are naturally disrupted via strong encounters in the cluster.

The numerical setup and initial properties for these simulations are similar to the work presented in (Chatterjee et al. 2013) which show that these models at the end of the simulation (at cluster age $t_{cl} \approx 12$ Gyr) attain properties including the core radius (r_c), the half-mass radius (r_h), central density (ρ_c), total mass (M_{cl}), and relaxation timescale at half-mass (t_{rh}) very similar to those observed in the GGCs. The only difference between these simulations and those presented in Chatterjee et al. (2013) is that models with zero initial f_b are discarded in this study since observations suggest that this is not the case for the GGCs in reality (Davis et al. 2008; Milone et al. 2012). In addition, the models presented in this study explores a larger range in initial f_b , and a slightly larger range in initial King concentration parameter W_0 . Using these highly realistic models we focus on understanding and characterizing the BSSs in globular clusters. For a detailed comparison of model properties with the observed properties of the GGCs readers are referred to Chatterjee et al. (2013).

2.1. Definition of blue straggler stars in our models

We define BSSs as stars that are still in the Hydrogen burning stage but with masses $> 1.1 M_{TO}$, where, M_{TO} is the main-sequence turn-off mass of the cluster. Note that in real clusters BSSs are chosen from a HRD as stars that are brighter and hotter than the MS turn-off. Unfortunately, the defining box for the BSSs vary from cluster to cluster depending on, for example, the quality of the data, and the scatter width of the MS (e.g., Leigh et al. 2007). Often the criterion used by the observers is that no MS star should be counted as a BSS and to ascertain that some actual BSSs can be discarded because of their proximity to the MS in the HRD. Thus it is hard to formulate a self-consistent luminosity and temperature based criterion that would be valid for all clusters. The mass-based criterion is much simpler and self-consistent to use for the models (although, inaccessible to observers). We find that in our models the BSSs identified using the mass-based criterion populate the expected region of synthetic HRDs created using our models (Figure 1 for an example). Since in this study we restrict ourselves to the theoretical understanding of the BSSs, in particular to understand the dominant formation channels and typical ages since formation, we simply adopt the mass-based definition for BSSs.

Due to the differences between the BSS identification criteria in our models and for observed clusters we find that although our models populate similar ranges in global properties such as r_c with the observed GGCs, we can identify more

BSSs than observed. For example, Figure 2 shows the number of BSSs within the core as a function of the core radius both for our models and for observed GGCs. The observed BSS counts are taken from the compilation in Leigh et al. (2007). Our models clearly have very similar core radii compared to the observed GGCs, however, we identify a larger number of BSSs in the core. This is expected since due to the uncertainty in determining the main-sequence turn-off in real clusters a significant number of BSSs can remain unidentified due to their proximity to the MS. Indeed, as part of a future study we have seen that if a more observationally motivated selection criteria for our model BSSs is used following the prescriptions of Leigh et al. (2007), the number of BSSs match well with the observations. In addition, the numbers of BSSs using either criteria are well correlated (Sills et al. 2013, in preparation). In addition to the differences in the identification criteria for the BSSs in our models with the observed, one should also note that within BSE, the rejuvenation prescription used for mergers, collisions and binary mass transfers assumes “full mixing” (for details see Hurley et al. 2000, 2002). This is a simplification. In reality, the amount of Hydrogen mixing and thus the degree of rejuvenation of the MS star depends on the kinematic details of the encounter (e.g., Sills et al. 1997, 2001). Because of this simplification the lifetimes of the BSSs after rejuvenation in our models are actually *upper limits* for non-rotating stars. This also leads to an over-production of BSSs in our models to some degree. This issue is further discussed in Section 5.

2.2. Classification of blue-straggler stars based on their formation channels

At the last snapshot of the model (t_{cl}) we identify the BSSs following the criterion described in Section 2. The full dynamical history for each of these BSSs is retrieved. We identify the interaction (such as SS collision, BS collision, or MTB) that had increased the mass of this star for the *first time* such that it would be identified as a BSS at t_{cl} . We classify each BSS into one of 5 groups according to this first interaction. The 5 groups are: stable mass transfer in a binary denoted as MTB, binary-stellar-evolution-driven merger/disruption denoted as SE, single-single collisions denoted as SS, collisions mediated by binary-single interactions denoted as BS, and collisions mediated by binary-binary encounters denoted as BB. Note that some BSSs may have a complicated history. For example, after a mass transfer episode in a binary, a MS star may have attained a BSS mass; later on this star could still collide with another MS star, further growing in mass and getting rejuvenated for a second time. For BSSs that are formed via such multiple channels, *only the first one* is considered for the classification to avoid ambiguity. In this particular example the formation channel for this BSS would be classified as MTB. Nevertheless, such complicated multiple episodes of rejuvenation of the same MS star are not frequent. There is another source of complication. A BSS classified as MTB (created via MTB) may have had interacted dynamically prior to the Roche-lobe overflow that had not changed its mass. Such an interaction would change the orbital properties of the binary, sending it along a very different evolutionary pathway that led to the mass transfer event that ultimately created the BSS. These perturbed MTB or SE channels are duly noted and discussed. However, interactions of this type do not change the formation classification of the BSSs.




FIG. 1.— Example of a synthetic Hertzsprung-Russell diagram for a representative model with many blue straggler stars (run110, Table 1). Dots denote all stars in the cluster. Red squares and blue triangles denote blue-straggler stars in single and binary systems, respectively at the integration stopping time of $t_{\text{cl}} = 12$ Gyr. Blue straggler stars are defined as H-burning stars with stellar masses $\geq 1.1 M_{\text{TO}}$, where M_{TO} is the main-sequence turn-off mass. BSSs in this model populate the expected region in the synthetic HRD. In addition, the single and binary main-sequences, giant branch, white-dwarf cooling sequence are clearly visible.

3. RADIAL DISTRIBUTION OF THE BLUE STRAGGLER STARS

One of the observationally accessible properties of the BSSs in the GGCs is their radial distribution. The radial distribution of the BSSs in a cluster depends on a collection of the host cluster's properties including ρ_c , relaxation time, and f_b . Thus analysis of the radial distribution of the BSSs has been proposed as a tool to observationally constrain the dominant formation channel, and various dynamical properties of the host GGCs. As a result, the radial distribution of the BSSs has been studied with interest for many GGCs via observations (e.g., Fusi Pecci et al. 1992; Dalessandro et al. 2008b; Ferraro et al. 1995; Beccari et al. 2008; Dalessandro et al. 2008a; Moretti et al. 2008a,b; Ferraro et al. 2012) as well as via theoretical modeling (e.g., Mapelli et al. 2006). Three qualitatively different types of radial distributions are observed among the GGCs, namely, single-peaked, double-peaked, and flat distributions (for a compilation see Ferraro et al. 2012).

To find the relative frequency of the BSSs in the cluster as a function of their radial positions we define a double-normalized BSS frequency R_{BSS} following Ferraro et al. (2012) given by

$$R_{\text{BSS}} = \frac{\delta N_{\text{BSS}}/N_{\text{BSS}}}{\delta L/L_{\text{cl}}}, \quad (1)$$

where δN_{BSS} denotes the number of BSSs within some radial bin, N_{BSS} denotes the total number of BSSs, δL denotes the stellar luminosity within that bin, and L_{cl} denotes the total luminosity of the cluster. R_{BSS} is a measure of whether the BSSs follow the luminosity density profile of the cluster or they are over/under abundant in some parts.

Figure 3 shows R_{BSS} as a function of the radial position in units of the core radius. As in some observed GGCs (e.g., M55) we find examples of double-peaked distributions (bottom panel). We also find examples of single-peaked distributions (top panel; e.g., observed in M80, Ferraro et al. 2012, and references therein). In our models we do not find flat radial distributions observed, for example, in NGC 2419 (Dalessandro et al. 2008b). This is expected since all our models are relaxed and hence a non-segregated radial distribution for the BSSs in our models is not expected. Ferraro et al. (2012) suggest that the minimum distance r_{min} from the center where the central peak of R_{BSS} ends in units of r_c is anti-correlated with the core relaxation timescale due to mass-segregation effects of the relatively high-mass BSSs. Results from our models are consistent with this finding. However, we find that the determination of r_{min} can have large errors and depends strongly on the radial bins used to calculate R_{BSS} .

4. RELATIVE IMPORTANCE OF BLUE STRAGGLER FORMATION CHANNELS




FIG. 2.— Number of BSSs in the core ($N_{\text{BSS},c}$) vs r_c . Black circles and red squares denote model and observed BSSs, respectively. The observed BSS numbers in the core are obtained from Leigh et al. (2007). Our models populate very similar r_c values compared to the observed GGCs. However, our models show higher $N_{\text{BSS},c}$ numbers due to the mass-based selection criteria. In an observed cluster many of these BSSs will not be identified as BSSs due to scatter in the main-sequence and the MS turn-off.

In this section we investigate the relative contributions from various BSS formation channels. Figure 4 shows an example of the branching ratio for the BSSs in the whole cluster and in the core for model run32 (Table 1). This is a model with a moderate central density $\rho_c = 7 \times 10^3 \text{ M}_\odot \text{ pc}^{-3}$. Collisions produce most of the BSSs in this model. Among the collisional BSSs channel SS contributes the least. The binary mediated collisional channels, namely BS and BB contribute about 36% and 50%, respectively. BSSs produced via the channel MTB contributes about 11% and SE contributes only 3.2%. As expected collisional channels dominate even more in this cluster if only core BSSs are considered. Another interesting aspect is that in this particular example model although the overall contribution from collisions via BS and BB channels increase if only core BSSs are considered compared to all BSSs in the cluster, contribution from the SS channel decreases. This trend is seen often in our models. This can be counter-intuitive to some degree since due to the higher central stellar densities, all forms of stellar collisions are expected to be more frequent within the core. This is because of the low density of single MS stars within the core at late times due to mass segregation.

About 60% of all BSSs formed via MTB or SE in this example model have had at least one prior strong encounter which did not make it a BSS. Thus for this particular model a significant fraction of even the MTB and SE BSSs have been dynam-

ically modified before binary stellar evolution could rejuvenate them. It will be very interesting to investigate in a future study whether such dynamical encounters typically enhance the possibility of MTB and SE BSS production relative to the field or inhibit it.

Branching ratios for various formation channels depend directly on the global properties, including the central density ρ_c of the host cluster. The higher the central density of a cluster, higher is the chance of a collisional formation of BSSs. Figure 5 shows the branching ratios for SS, BS, BB, MTB, and SE formation channels as a function of ρ_c . For models with central densities $\rho_c > \text{few} \times 10^3 \text{ M}_\odot \text{ pc}^{-3}$ collisional channels clearly dominate. In lower density clusters contribution from MTB increases. Throughout the full range in binary fraction in our models ($f_b = 0.05 - 0.30$), the same trend is observed. This indicates, that even if f_b is increased (within the studied range in ρ_c) the relative contribution from MTB does not increase proportionally. While increasing f_b increases the probability that a BSS can be created via MTB (simply because there are more binaries), the increased number of binaries also increases the BS and BB BSS formation rates. Typical ρ_c values in the GGCs are orders of magnitude higher than $\sim 10^3 \text{ M}_\odot \text{ pc}^{-3}$ (e.g., Harris 1996; Chatterjee et al. 2010, 2013). Hence, we conclude that the collisional channels dominate BSS production for the populations observed in majority of the GGCs. Only about 30% of all GGCs have observed




FIG. 3.— Examples of two qualitatively different radial distributions for our model BSSs. Vertical and horizontal axes denote the double-normalized BSS frequency R_{BSS} (Equation 1) and radial position in units of r_c , respectively. Top panel shows an example of centrally peaked radial distribution (run108; Table 1). Bottom panel shows an example of a double peaked distribution (run110) one near the center $r/r_c < 5$ and a second peak at large $r/r_c \approx 22$. A zone of avoidance is observed at r/r_c between about 5 and 17.




FIG. 4.— Example of branching ratios for the various formation channels of BSSs at $t = 12$ Gyr (run32; Table 1). This model has a moderate central density ($\rho_c \approx 7 \times 10^3 M_\odot \text{pc}^{-3}$; Table 1). This particular model was chosen as an example due to its high number (312) of BSSs. The dominant formation channel is stellar collisions in this model. Among the collisional BSS production channels collisions mediated by BB interactions dominate followed by BS mediated collisions. SS collisions are rare. Mass transfer in a binary is the dominant channel among the binary-stellar-evolution driven BSS production channels. Binary stellar evolution driven mergers or disruptions contribute to a small degree.




FIG. 5.— Branching ratios of various formation channels for all BSSs in our models vs the respective cluster's ρ_c . The circles, squares, and triangles denote collisional, MTB, and SE channels, respectively. Clearly, for $\rho_c > \text{few} \times 10^3 M_\odot \text{pc}^{-3}$ collisional formations channels dominate. SE mergers are rare throughout the full range in ρ_c in our models. Top to bottom models with initial $f_b = 0.05, 0.1, 0.2$, and 0.3 are presented, respectively.




FIG. 6.— Same as Figure 5 but including only the BSSs within the core. As expected collisional channels dominate the core BSSs. SE mergers are rare throughout the range of ρ_c in our models. MTB channel only becomes comparable to collisional channels if $\rho_c < 10^3 M_\odot \text{pc}^{-3}$.

ρ_c values consistent to have BSS populations predominantly from MTB (Harris 1996). If only the core population of BSSs is considered, the importance of the collisional channels increases as expected. We find that throughout the full range of central densities in our models collisional channels dominate the formation of core BSSs (Figure 6).

Although binary stellar evolution is not the dominant channel for a large range of central densities typical of the GGCs, binaries themselves are nevertheless crucial for significant BSS production. Figure 7 shows the relative contribution between various collisional formation channels of the BSSs as a function of ρ_c . For the full range in f_b in our models BSSs produced via SS are rare. Binary-mediated collisions (BS or BB) dominate. As f_b increases, so does the relative contribution from BB. For example, for models with initial $f_b = 0.05$, the BS channel contributes more to the collisional BSSs compared to the BB channel for all ρ_c values. In contrast, for $f_b = 0.3$ the contribution from the BB channel dominates for all ρ_c values we have considered.

Figure 9 shows the fraction of BSSs produced via binary stellar evolution (MTB and SE collectively) that have been dynamically perturbed before the mass transfer process has occurred. These interactions can potentially alter the orbital properties of these binaries initiating very different evolutionary pathways that lead to mass transfer. BSSs formed from these disturbed binaries clearly would not have formed the BSSs with the exact same properties and at the same times without these encounters. As expected, for the models with

higher ρ_c values MTB and SE BSSs have a higher chance of being perturbed before production. For a given ρ_c a range of these values are also seen. This is due to the different concentration values. Models with initially lower concentration leads to a smaller fraction of perturbed MTB and SE BSSs. The majority of the models with $\rho_c > 10^4 M_\odot \text{pc}^{-3}$ produce more perturbed MTB and SE BSSs than unperturbed. Low density ($\rho_c < 10^3$) models on the other hand will have a much larger fraction of BSSs via the MTB and SE channels created from the primordial undisturbed binaries. Note that this is in direct contrast from some earlier observational studies, e.g., by Sollima et al. (2008, further discussion in Section 6).

5. TIME SINCE FORMATION FOR THE BLUE STRAGGLER STARS

How long a BSS will live after the encounter that formed and possibly rejuvenated it depends on the supply of new Hydrogen that can be mixed into its core for burning. The degree of rejuvenation depends on the detailed kinematics of the encounter (e.g., Sills et al. 1997, 2001) as well as the stellar properties such as rotation, and the ages of the parent stars prior to interaction (Chen & Han 2009). Incorporating these details is hard and beyond the scope of this study. Instead we adopt a full mixing prescription used in the stellar evolution code BSE (Hurley et al. 2000, 2002) that has been used in this study as part of our cluster evolution code CMC and is widely used in most N -body codes. It has been previously found that this full-mixing prescription adopted in BSE predicts a longer rejuvenated lifetime than found in detailed




FIG. 7.— Branching ratios between various collisional BSS formation channels vs ρ_c . Squares, triangles, and circles denote BS, BB, and SS collisions, respectively. Throughout the studied range of ρ_c and f_b , single-single collisions are rare compared to binary-single and binary-binary collision channels. At high $f_b = 0.3$ values binary-binary collisions dominate over binary-single collisions for the production of BSSs.

stellar evolution models assuming head-on collisions (e.g., Glebbeek et al. 2008; Glebbeek & Pols 2008). Hence the time since formation for the BSSs in these models (especially created via head-on collisions) should be treated as upper limits.

We define the elapsed times since formation $t_{\text{age}} \equiv t_{\text{cl}} - t_{\text{form}}$, where t_{form} is the time when for the first time a star achieved a mass that will give it a BSS status at the final cluster age t_{cl} via some channel. Recent detailed numerical studies indicate that most GGCs (observationally categorized as the non-core-collapsed clusters) are still contracting (e.g., Chatterjee et al. 2013). Hence, relevant global cluster properties such as ρ_c , r_c , and r_h do not attain steady-state values typical of, e.g., core-collapsed a.k.a. binary-burning clusters. A priori, a BSS can have any t_{age} between zero and the cluster’s age. However, there may also be a “typical” t_{age} corresponding to a peak in the age distribution. Thus to understand the environment that had produced the present-day BSSs one must take into account the evolving properties of the cluster for at least this typical t_{age} . Unfortunately, t_{age} is impossible to determine purely from observations and thus results from detailed cluster models is the only way.

Figure 10 shows examples of the distributions of t_{age} in our models differing by almost two orders of magnitude in their ρ_c values. The BSSs in the high-density model are typically younger compared to those in the low-density model. The broadly peaked distributions seen here are qualitatively similar in all our models. However, the exact positions of

these peaks depend on the cluster properties. For the low-density model the bulk of the BSS population have ages between about $t_{\text{age}} \approx 0 - 5$ Gyr. This is consistent with previous results suggested for the observed BSS population in 47 Tuc (Sills et al. 2000). In contrast, the BSSs in the high-density model typically have t_{age} between 0–3 Gyr. In each case there is a long tail in the t_{age} distribution extending to about 6 Gyr. Half of all BSSs in the low-density model is contained within an age of 2 Gyr, whereas, for the high-density model this value is 0.9 Gyr.

The difference in the peaks of the age distributions between the high and low-density models is expected. In a high-density model the dominant formation channel for BSSs is stellar collisions (SS, BS, and BB; Figures 4, 5). In contrast MTB is the dominant formation channel for sufficiently low density ($\rho_c \leq 10^3 M_\odot \text{pc}^{-3}$) clusters (Figure 5). Stellar collisions typically form higher mass BSSs than those formed via MTB. Since a lower mass BSS has a longer residual lifetime than a higher mass counterpart, the typical t_{age} becomes shorter as ρ_c increases due to higher contribution from collisional BSS-formation channels. Indeed we find that the median for the t_{age} values for the BSSs in our models negatively correlate with ρ_c (Figure 11). Due to the same reason the median value of t_{age} for the core BSSs is typically slightly lower compared to all BSSs. This effect is more prominent for higher f_b models. The number of BSSs outside r_c is low for low f_b models resulting in little difference between the




FIG. 8.— Same as Figure 7 but including only the BSSs within the core.

median t_{age} values for all BSSs in the clusters and for the BSSs inside r_c only. We find that the oldest BSSs in a cluster can be almost as old as the cluster itself.

6. SPECIFIC NUMBER OF BLUE-STRAGGLER STARS VS BINARY FRACTION AND Γ

After the large-scale surveys using the Hubble Space Telescope (e.g., the ACS survey; Sarajedini et al. 2007; Dotter et al. 2007; Milone et al. 2008) provided high resolution observations for the GGCs, a large number of studies have investigated various correlations between the number of BSSs in these clusters and their global properties (e.g., Leigh et al. 2007; Sollima et al. 2008; Knigge et al. 2009; Leigh et al. 2013). In particular, the correlation or lack thereof between the core BSSs ($N_{\text{BSS},c}$) and two cluster properties namely the binary fraction in the core ($f_{b,c}$) and the collisional Γ have been studied in much detail. Both of these quantities are of great dynamical importance. The presence of primordial binaries increases the chance of all BSS formation channels except SS. Whereas, BSS formation via SS is directly related to the collisional Γ given by

$$\Gamma = \left(\frac{\rho_c}{M_{\odot} \text{pc}^{-3}} \right)^2 \left(\frac{r_c}{\text{pc}} \right)^3 \left(\frac{v_{c,\sigma}}{\text{km s}^{-1}} \right)^{-1} \quad (2)$$

(Pooley & Hut 2006). Especially, after it was found that the number of low-mass X-ray binaries and CVs in a cluster is strongly correlated with Γ (e.g., Pooley & Hut 2006) the same correlation was sought for the BSSs in the core

since all of them are dynamically created populations. However, it was found that the specific number of BSSs in the core, normalized by another population similar in mass to account for the fact that a higher- N clusters should naturally have a higher number of BSSs, e.g., the horizontal branch stars, $N_{\text{BSS},c}/N_{\text{HB},c}$, is rather insensitive to the estimated Γ . In contrast, it is found that $N_{\text{BSS},c}/N_{\text{HB},c}$ shows clear correlations with the core binary fraction in the GGCs (e.g., Leigh et al. 2007; Sollima et al. 2008; Knigge et al. 2009; Leigh et al. 2013). These two pieces of information led to the claim that most of the BSSs in the GGCs come via binary stellar evolution channels (MTB and SE) and collisional channels (SS, BS, and BB) are unimportant (e.g., Sollima et al. 2008; Knigge et al. 2009). However, our models suggest that collisions dominate BSS production in majority of the GGCs. We now investigate further this apparent discrepancy.

Figure 12 shows the specific number of BSSs ($N_{\text{BSS}}/N_{\text{HB}}$) in our models as a function of the overall f_b . It also shows the specific number of BSSs in the core ($N_{\text{BSS},c}/N_{\text{HB},c}$) as a function of the core binary fraction $f_{b,c}$. Clearly, $N_{\text{BSS}}/N_{\text{HB}}$ is strongly correlated with f_b and $N_{\text{BSS},c}/N_{\text{HB},c}$ is strongly correlated with $f_{b,c}$. This is not at all surprising. Results in Section 4 clearly shows that throughout the studied range in ρ_c the dominant BSS production channel is binary-mediated collisions (BS and BB). Hence, as f_b increases so does the number of BSSs. Although, naturally the number of BSSs from the MTB and SE channels also increases with increasing f_b , their relative importance is always low compared to




FIG. 9.— Fraction of perturbed BSSs created via MTB or SE mergers as a function of ρ_c . Only perturbations before the production of the BSSs are taken into account. This fraction is directly related to the fraction of binary stellar evolution driven BSSs (MTB and SE) that would have not been produced with the same properties had there not been a previous encounter. In addition, once produced, these BSSs may interact further which was not taken into account. Thus, these fractions indicate a lower limit for the fraction of observed BSSs that have not been dynamically altered. Black circles and red squares denote all BSSs in the models and only those within the core, respectively. Each panel from top to bottom show initial $f_b = 0.05, 0.1, 0.2$, and 0.3 , respectively.

BS and BB channels for all $\rho_c > 10^3 M_\odot \text{pc}^{-3}$. Thus the observed correlation between $N_{\text{BSS},c}/N_{\text{HB},c}$ with $f_{b,c}$ does not actually indicate a predominantly MTB BSS formation, rather that most collisions are binary-mediated.

Figure 13 shows $N_{\text{BSS},c}/N_{\text{HB},c}$ as a function of Γ calculated using true cluster values in the code. We find a positive correlation between the $N_{\text{BSS},c}/N_{\text{HB},c}$ and the theoretical Γ values. This is only expected since our results suggest that the dominant formation channel for the BSSs in typical GGCs is collisions of some kind. This appears to be in disagreement with the observed trend. Closer inspection reveals that the apparent discrepancy is likely coming from the differences in theoretically calculated Γ values and those that are estimated from observed clusters. Note that the Γ values are calculated using three cluster properties, r_c , ρ_c , and the velocity dispersion ($v_{c,\sigma}$) in the core. Most recently Chatterjee et al. (2013) show that it is important to be careful about the definitions of all of the above quantities while comparing theoretical models with observed. To find what an observer would estimate for the Γ in our model clusters (Γ_{obs} to differentiate from true Γ) we estimate an observed r_c as prescribed in Chatterjee et al. (2013), measure the central density as an observer would from the surface brightness profile, and calculate $v_{c,\sigma}$ taking into account only the bright MS ($M_* > 0.3 M_\odot$) and giant stars in the cluster. Using these “observed” values for r_c , ρ_c , and $v_{c,\sigma}$ we then calculate the observed value of Γ_{obs} for all our mod-

els. We find that the theoretical Γ values are not retrieved easily from Γ_{obs} (Figure 14) indicating that the Γ_{obs} values estimated in observed GGCs are not reliable indicators of the actual collision rates. Figure 15 shows $N_{\text{BSS},c}/N_{\text{HB},c}$ as a function of Γ_{obs} for all our models. We find that the correlation between $N_{\text{BSS},c}/N_{\text{HB},c}$ and Γ as seen in Figure 13 is not apparent between $N_{\text{BSS},c}/N_{\text{HB},c}$ and Γ_{obs} . Thus our models suggest that although most BSSs are created via binary-mediated collisional channels in typical GGCs, $N_{\text{BSS},c}/N_{\text{HB},c}$ may appear insensitive to the observed estimates of Γ_{obs} .

7. CONCLUSIONS

We have identified the BSSs from a large ensemble of realistic simulated clusters using a simple and well defined mass-based criterion. The BSSs populate the expected region in a synthetic HRD (Figure 1). Although the stellar properties and numbers of BSSs in various clusters can be determined observationally, it is hard to estimate the typical t_{age} for observed BSSs. Moreover, the formation channels of these observed BSSs are also unknown. Our detailed realistic N -body models provide, for the first time, rigorous constraints on the branching ratios between various formation channels, distributions of t_{age} and also shed light on the observed correlations between the number of BSSs in a cluster and its dynamical properties including f_b , and Γ .

We have identified the relative importance of the various




FIG. 10.— Examples for distributions of the time elapsed since formation (t_{age}) for the BSSs for two representative models, one for low (run61) and another for high ρ_c (run113; Table 1) with central densities varying by two orders of magnitude. The solid (black) and dashed (blue) lines represent the low and high density clusters, respectively. The top and bottom panels are for all BSSs and those within r_c , respectively. For MTB BSSs the formation time t_{form} is set at the beginning of Roche-lobe overflow. The BSSs in the low-density model shows a larger age-spread compared to those in the high-density model. Low-density models have a larger relative contribution from MTB channel. Since MTB BSSs are typically less massive compared to those created via collisions, the residual lifetime for the MTB BSSs is longer than those produced via collisions. Throughout the studied range in ρ_c , our models indicate that a significant fraction of today’s observed BSSs had been created many Gyrs ago.

BSS formation channels. In particular we divide all BSSs in our models in 5 different categories based on their detailed dynamical history. We find that collisions play a major role in BSS formation for a large range of central densities typical of globular clusters ($\rho_c > 10^3 M_\odot \text{pc}^{-3}$; Figures 4, 5, and 6). BSSs in lower density clusters ($\rho_c < 10^3 M_\odot \text{pc}^{-3}$) can have a significant contribution from the MTB channel (up to $\approx 60\%$ in some of our models). This also suggests that for open cluster densities (typically $< 10^2 M_\odot \text{pc}^{-3}$) the dominant channel of BSS formation should be MTB. This picture can only change if in reality the degree of mixing from the MTB channel for some reason is very different than that from collisions. One extreme case is, for example, collisions lead to minimal mixing and no Hydrogen can be supplied to the core. If this is true then there would be no rejuvenation from collisions and thus, collisions would produce BSSs that live in the BSS stage only a short time. If, however, degree of rejuvenation from collisional channels and MTB is similar, then our results remain valid.

The above finding is of great interest. It has been seen in the observed BSS populations in the Galaxy that the number of BSSs in the core does not show a positive correlation with the collisional Γ parameter (Davies et al. 2004). It has also been noted that there is a strong correlation between the

number of core BSSs and the binary fraction in the cluster (Knigge et al. 2009). These results had led to conclusions in some studies that MTB from undisturbed primordial binaries is the dominant formation channel of BSSs (Sollima et al. 2008). Our results contradict this conclusion and suggest that binary mediated collisions is the dominant BSS formation channel instead. Although, MTB is not the major contributor in BSS formation in typical globular clusters, binaries play a crucial role in BSS formation. Because of the higher interaction cross-section of the binaries they are much more likely to interact compared to the single stars. Moreover, single MS stars tend to reside outside the core at late times due to mass-segregation. For typical globular cluster velocity dispersions the branching ratio for collisions in binary interactions is significant (Fregeau et al. 2003, 2004). BS and BB interactions lead to the majority of the collisions in globular clusters creating the BSSs (Figure 7, and 8). Hence, we concur with the suggestion that binaries are of great importance in creating BSSs in a cluster. However, from our models we conclude that this is not because most BSSs are created from MTB. The apparent lack of correlation between the specific BSS number and the collisional Γ parameter had been noted as suggestive to a predominantly non-collisional channel of BSS formation previously (e.g., Sollima et al. 2008).




FIG. 11.— The median values for t_{age} of the BSSs in our models vs ρ_c . Black circles and red squares denote all BSSs and BSSs in the core, respectively. Top to bottom the panels denote models with initial $f_b = 0.05, 0.1, 0.2$, and 0.3 , respectively. The median t_{age} values for all initial f_b show an anti-correlation with ρ_c . The denser the core, the higher is the contribution of the collisional channels towards BSS production and the lower is the typical median age of the BSSs.

Our models suggest that this is clearly *not* the case. Even for the BSSs created via MTB or SE channels, a large fraction of them have been dynamically perturbed especially for clusters with $\rho_c > 10^3 M_\odot \text{pc}^{-3}$. Close inspection of our models reveals that although $N_{\text{BSS},c}/N_{\text{HB},c}$ do correlate weakly with the actual theoretically calculated Γ in our models, they do not with the Γ_{obs} estimated taking into account the observational difficulties in measuring this quantity.

We also determined upper limits for the typical t_{age} for the BSSs in our models. In our models the oldest BSSs can be almost as old as the cluster itself. The typical ages of the BSSs are, however, between 1 – 5 Gyr (depending on ρ_c ; Figure 10, 11 and Table 1). Thus, present-day BSSs are possibly affected by the dynamical history of their host clusters during the last several billion years. However, we caution that due to the assumption of full-mixing inherent in BSE (Hurley et al. 2000, 2002), the stellar evolution routine used in our code CMC, the t_{age} values should be treated as upper limits, in particular for non-rotating stars (e.g., Glebbeek & Pols 2008).

Throughout this study we have defined BSSs using a simple mass-based criterion that is easy in theory but inaccessible in real observations. Although, the BSSs in our models identified based on this straight-forward criterion populate the expect regions in the HRD, we note that a fraction of the theoretically identified BSSs in these models would not have

been identified as BSSs based on a more observationally motivated criteria described, for example, in Leigh et al. (2007). We have started investigating the properties of the BSSs in our models based on this observationally motivated criteria as part of another ongoing study. The theoretically extracted BSSs in our models is a superset of the BSSs that would be extracted using the observationally motivated criterion.

8. ACKNOWLEDGEMENTS

Support for this study was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Incorporated, under NASA contract NAS5-26555. The grant identifying number is HST-AR-12829.01-A. SC also acknowledges support from the Department of Astronomy, University of Florida. FAR acknowledges support from NASA ATP Grant NNX09AO36G at Northwestern University. AS is supported by the Natural Sciences and Engineering Research Council of Canada. EG is supported by NWO under grant 639.041.129. All author would like to thank the Kavli Institute for Theoretical Physics for their hospitality, where this project began. This work was supported in part by the National Science Foundation under Grant No. NSF PHY11-25915.

REFERENCES

- Beccari, G., Lanzoni, B., Ferraro, F. R., Pulone, L., Bellazzini, M., Fusi Pecci, F., Rood, R. T., Giallongo, E., Ragazzoni, R., Grazian, A., Baruffolo, A., Bouche, N., Buschkamp, P., De Santis, C., Diolaiti, E., & et al. 2008, *ApJ*, 679, 712
- Chatterjee, S., Fregeau, J. M., Umbreit, S., & Rasio, F. A. 2010, *ApJ*, 719, 915




FIG. 12.— Left Panel: The ratio of the number of BSSs (N_{BSS}) to the number of horizontal-branch stars (N_{HB}) vs the final f_b . Right Panel: The ratio of the number of core BSSs ($N_{\text{BSS},c}$) to the number of core horizontal-branch stars ($N_{\text{HB},c}$) vs the final core binary fraction ($f_{b,c}$). The specific BSS frequencies $N_{\text{BSS}}/N_{\text{HB}}$ and $N_{\text{BSS},c}/N_{\text{HB},c}$ are strongly correlated with f_b and $f_{b,c}$, respectively. Although the dominant BSS production channel for these models is stellar collisions, most of these collisions are binary mediated. Hence, a higher binary fraction leads to higher number of collisions and thus a higher number of BSSs if all else is kept unchanged.

Chatterjee, S., Umbreit, S., Fregeau, J. M., & Rasio, F. A. 2013, *MNRAS*, 429, 2881
 Chen, X. & Han, Z. 2009, *MNRAS*, 395, 1822
 Dalessandro, E., Lanzoni, B., Ferraro, F. R., Rood, R. T., Milone, A., Piotto, G., & Valenti, E. 2008a, *ApJ*, 677, 1069
 Dalessandro, E., Lanzoni, B., Ferraro, F. R., Vespe, F., Bellazzini, M., & Rood, R. T. 2008b, *ApJ*, 681, 311
 Davies, M. B., Piotto, G., & de Angeli, F. 2004, *MNRAS*, 349, 129
 Davis, D. S., Richer, H. B., Anderson, J., Brewer, J., Hurley, J., Kalirai, J. S., Rich, R. M., & Stetson, P. B. 2008, *AJ*, 135, 2155
 Dotter, A., Chaboyer, B., Jevremović, D., Baron, E., Ferguson, J. W., Sarajedini, A., & Anderson, J. 2007, *AJ*, 134, 376
 Ferraro, F. R., Fusi Pecci, F., & Bellazzini, M. 1995, *A&A*, 294, 80
 Ferraro, F. R., Lanzoni, B., Dalessandro, E., Beccari, G., Pasquato, M., Miocchi, P., Rood, R. T., Sigurdsson, S., Sills, A., Vesperini, E., Mapelli, M., Contreras, R., Sanna, N., & Mucciarelli, A. 2012, *Nature*, 492, 393
 Fregeau, J. M., Cheung, P., Portegies Zwart, S. F., & Rasio, F. A. 2004, *MNRAS*, 352, 1
 Fregeau, J. M., Gürkan, M. A., Joshi, K. J., & Rasio, F. A. 2003, *ApJ*, 593, 772
 Fregeau, J. M. & Rasio, F. A. 2007, *ApJ*, 658, 1047
 Fusi Pecci, F., Ferraro, F. R., Corsi, C. E., Cacciari, C., & Buonanno, R. 1992, *AJ*, 104, 1831
 Glebbeek, E. & Pols, O. R. 2008, *A&A*, 488, 1017
 Glebbeek, E., Pols, O. R., & Hurley, J. R. 2008, *A&A*, 488, 1007
 Harris, W. E. 1996, *AJ*, 112, 1487
 Hurley, J. R., Pols, O. R., & Tout, C. A. 2000, *MNRAS*, 315, 543
 Hurley, J. R., Tout, C. A., & Pols, O. R. 2002, *MNRAS*, 329, 897
 Joshi, K. J., Nave, C. P., & Rasio, F. A. 2001, *ApJ*, 550, 691
 Joshi, K. J., Rasio, F. A., & Portegies Zwart, S. 2000, *ApJ*, 540, 969
 Knigge, C., Leigh, N., & Sills, A. 2009, *Nature*, 457, 288

Kozai, Y. 1962, *AJ*, 67, 591
 Kroupa, P. 2001, *MNRAS*, 322, 231
 Leigh, N., Knigge, C., Sills, A., Perets, H. B., Sarajedini, A., & Glebbeek, E. 2013, *MNRAS*, 428, 897
 Leigh, N., Sills, A., & Knigge, C. 2007, *ApJ*, 661, 210
 Lombardi, Jr., J. C., Rasio, F. A., & Shapiro, S. L. 1995, *ApJ*, 445, L117
 —. 1996, *ApJ*, 468, 797
 Lombardi, Jr., J. C., Warren, J. S., Rasio, F. A., Sills, A., & Warren, A. R. 2002, *ApJ*, 568, 939
 Mapelli, M., Sigurdsson, S., Ferraro, F. R., Colpi, M., Possenti, A., & Lanzoni, B. 2006, *MNRAS*, 373, 361
 Mathieu, R. D. & Geller, A. M. 2009, *Nature*, 462, 1032
 Milone, A. P., Bedin, L. R., Piotto, G., Anderson, J., King, I. R., Sarajedini, A., Dotter, A., Chaboyer, B., Marin-Franch, A., Majewski, S., Aparicio, A., Hempel, M., Paust, N. E. Q., Reid, I. N., Rosenberg, A., & Siegel, M. 2008, *ApJ*, 673, 241
 Milone, A. P., Piotto, G., Bedin, L. R., Aparicio, A., Anderson, J., Sarajedini, A., Marino, A. F., Moretti, A., Davies, M. B., Chaboyer, B., Dotter, A., Hempel, M., Marin-Franch, A., Majewski, S., Paust, N. E. Q., Reid, I. N., Rosenberg, A., & Siegel, M. 2012, *A&A*, 540, A16
 Moretti, A., de Angeli, F., & Piotto, G. 2008a, *VizieR Online Data Catalog*, 348, 30183
 —. 2008b, *A&A*, 483, 183
 Pattabiraman, B., Umbreit, S., Liao, W.-k., Choudhary, A., Kalogera, V., Memik, G., & Rasio, F. A. 2013, *ApJS*, 204, 15
 Perets, H. B. & Fabrycky, D. C. 2009, *ApJ*, 697, 1048
 Piotto, G., King, I. R., Djorgovski, S. G., Sosin, C., Zoccali, M., Saviane, I., de Angeli, F., Riello, M., Recio-Blanco, A., Rich, R. M., Meylan, G., & Renzini, A. 2002, *VizieR Online Data Catalog*, 339, 10945
 Pooley, D. & Hut, P. 2006, *ApJ*, 646, L143
 Sandage, A. R. 1953, *AJ*, 58, 61




FIG. 13.— Specific number of the BSSs in the core ($N_{BSS,c}/N_{HB,c}$) vs the collisional Γ parameter (Pooley & Hut 2006) theoretically calculated in our models (Equation 2). A weak positive correlation between $N_{BSS,c}/N_{HB,c}$ and Γ is found.

Sarajedini, A., Bedin, L. R., Chaboyer, B., Dotter, A., Siegel, M., Anderson, J., Aparicio, A., King, I., Majewski, S., Marín-Franch, A., Piotto, G., Reid, I. N., & Rosenberg, A. 2007, *AJ*, 133, 1658
 Sills, A., Bailyn, C. D., Edmonds, P. D., & Gilliland, R. L. 2000, *ApJ*, 535, 298
 Sills, A., Faber, J. A., Lombardi, Jr., J. C., Rasio, F. A., & Warren, A. R. 2001, *ApJ*, 548, 323

Sills, A., Lombardi, Jr., J. C., Bailyn, C. D., Demarque, P., Rasio, F. A., & Shapiro, S. L. 1997, *ApJ*, 487, 290
 Sollima, A., Lanzoni, B., Beccari, G., Ferraro, F. R., & Fusi Pecci, F. 2008, *A&A*, 481, 701
 Umbreit, S., Fregeau, J. M., Chatterjee, S., & Rasio, F. A. 2012, *ApJ*, 750, 31




FIG. 14.— Theoretically calculated Γ vs the Γ_{obs} values as an observer would estimate from our models. Γ and Γ_{obs} show no correlation. The lack of correlation between Γ_{obs} and Γ indicates that the canonical Γ_{obs} values measured in observed clusters may not be a reliable measure for the actual collisional rate in the GGCs. This is not surprising since Γ_{obs} depends on many structural parameters of a cluster (Equation 2) all of which can contribute errors in this estimate. We find that the estimate of $r_{c,\text{obs}}$ (e.g., Chatterjee et al. 2013) dominates the errors in estimating Γ_{obs} .




FIG. 15.— Same as Figure 13 but here Γ is calculated as an observer would if observing our model. The correlation observed in Figure 13 disappears.

TABLE 1
MODEL PROPERTIES AND DYNAMICAL PROPERTIES OF BSSs.

Name	t_{cl} (Gyr)	M ($10^4 M_{\odot}$)	f_b	$f_{b,c}$	ρ_c ($10^4 M_{\odot} \text{pc}^{-3}$)	r_c (pc)	N_{BSS}					$N_{BSS,c}$					t_{age} (Gyr)	
							Coll			MTB	SE	Coll			MTB	SE	Med	Max
							SS	BS	BB			SS	BS	BB				
run1	12	14	0.05	0.09	1.03	0.9	2	11	3	1	0	1	10	3	0	0	0.9	5.1
run2	12	21	0.05	0.09	0.74	1.2	4	24	4	2	1	3	15	2	0	0	1.0	5.8
run3	12	28	0.05	0.08	0.68	1.4	6	19	10	3	1	2	15	5	3	0	1.9	5.4
run4	12	36	0.05	0.07	0.67	1.5	11	30	7	10	0	7	17	5	5	0	1.6	5.0
run5	12	14	0.09	0.17	0.88	1.0	2	20	27	1	0	1	17	20	1	0	1.2	7.3
run6	12	22	0.09	0.16	0.68	1.3	4	24	19	8	1	1	14	10	4	0	1.6	5.4
run7	12	29	0.09	0.15	0.58	1.5	5	45	24	10	0	1	32	16	4	0	1.8	6.8
run8	12	36	0.09	0.14	0.58	1.6	10	38	25	15	1	5	25	18	6	1	1.9	7.3
run9	12	15	0.18	0.29	0.50	1.2	1	17	30	5	2	1	13	20	5	0	1.5	10.0
run10	12	23	0.18	0.29	0.61	1.3	1	49	73	15	3	0	37	46	7	1	1.2	5.4
run11	12	30	0.18	0.27	0.55	1.5	4	50	78	28	1	3	31	51	9	1	1.8	7.1
run12	12	38	0.18	0.25	0.50	1.7	3	75	60	44	1	2	52	40	18	0	1.5	5.9
run13	12	16	0.27	0.40	0.40	1.3	0	26	45	8	2	0	13	33	3	1	1.1	7.0
run14	12	24	0.27	0.39	0.46	1.5	3	44	99	17	4	0	25	61	8	1	1.2	9.1
run15	12	32	0.27	0.37	0.44	1.6	5	64	115	38	8	2	41	81	16	2	1.7	10.0
run16	12	40	0.27	0.36	0.45	1.8	2	97	106	49	4	1	63	76	19	2	1.7	12.0
run17	12	14	0.05	0.11	3.36	0.7	6	22	8	0	0	4	15	6	0	0	0.7	4.5
run18	12	21	0.05	0.09	1.66	1.0	2	14	9	1	1	0	5	9	0	1	1.5	6.3
run19	12	28	0.04	0.08	1.18	1.2	10	38	7	1	3	4	24	5	0	2	1.2	5.9
run20	12	36	0.04	0.08	1.15	1.3	12	36	12	2	2	4	25	8	1	1	1.1	7.1
run21	12	14	0.09	0.19	2.13	0.8	1	24	26	1	1	0	19	20	0	0	1.3	5.3
run22	12	22	0.09	0.16	1.20	1.1	5	49	29	1	2	1	33	22	1	1	1.0	6.0
run23	12	29	0.09	0.15	0.95	1.3	10	51	24	9	2	5	30	17	6	1	1.2	10.9
run24	12	36	0.09	0.15	0.98	1.3	10	61	35	8	2	6	41	23	2	2	1.6	9.2
run25	12	15	0.18	0.30	0.94	1.0	1	25	52	0	0	0	16	40	0	0	1.2	5.9
run26	12	23	0.18	0.29	0.91	1.2	2	53	61	13	1	0	36	41	4	1	1.4	11.5
run27	12	30	0.18	0.27	0.76	1.4	3	48	59	27	2	2	35	35	13	1	1.8	9.1
run28	12	38	0.18	0.26	0.82	1.4	2	83	110	28	3	0	57	83	12	1	1.4	6.4
run29	12	16	0.27	0.40	0.52	1.2	0	29	62	8	2	0	15	41	5	1	1.6	6.4
run30	12	24	0.27	0.39	0.55	1.4	0	58	98	11	6	0	35	71	4	0	1.4	6.9
run31	12	32	0.27	0.37	0.55	1.5	1	54	96	29	4	1	35	62	9	2	1.3	7.0
run32	12	40	0.27	0.36	0.70	1.5	1	112	156	33	10	0	66	107	11	1	1.7	8.6
run33	11	14	0.05	0.10	3.33	0.7	3	19	8	2	1	2	15	6	0	1	1.3	3.8
run34	11	21	0.05	0.10	3.11	0.8	6	24	10	2	0	3	16	9	2	0	0.8	3.9
run35	11	29	0.04	0.09	2.67	0.9	9	43	10	1	1	3	29	10	0	1	1.1	5.8
run36	11	36	0.04	0.08	2.20	1.0	12	39	13	1	3	6	28	8	0	3	1.1	4.6
run37	11	14	0.09	0.17	1.75	0.8	2	21	21	2	0	1	15	15	1	0	1.2	6.2
run38	11	22	0.09	0.18	2.26	0.9	2	47	19	1	1	1	30	11	0	0	0.9	5.5
run39	11	29	0.09	0.16	1.75	1.0	6	62	37	5	6	1	44	27	1	5	1.3	6.8
run40	11	37	0.09	0.15	1.88	1.1	7	60	27	12	3	3	42	23	7	2	1.5	9.1
run41	11	15	0.18	0.32	1.67	0.8	0	28	47	1	1	0	18	28	0	0	1.0	5.6
run42	11	23	0.18	0.31	1.85	1.0	5	61	74	8	1	1	38	50	5	0	1.0	7.1
run43	11	30	0.18	0.28	1.24	1.2	3	52	81	10	2	2	40	67	4	2	1.4	5.9
run44	12	38	0.18	0.27	1.55	1.2	6	85	80	12	4	5	62	60	4	3	1.2	7.1
run45	12	15	0.27	0.42	1.11	1.0	2	30	76	3	3	0	20	53	1	0	1.3	7.2
run46	11	24	0.27	0.40	1.04	1.2	2	53	103	10	7	1	33	74	4	3	1.2	6.0
run47	12	32	0.27	0.38	1.06	1.2	4	73	127	11	4	1	46	85	4	1	1.4	10.0
run48	12	40	0.27	0.37	1.05	1.3	3	77	161	26	4	2	53	115	6	2	1.3	8.1
run49	12	14	0.05	0.07	0.09	2.1	0	8	1	4	0	0	4	1	1	0	2.9	4.2
run50	12	21	0.05	0.07	0.11	2.3	1	9	3	7	1	1	5	2	5	1	2.2	6.5
run51	12	29	0.05	0.07	0.12	2.4	5	10	0	8	0	1	6	0	4	0	1.9	7.4
run52	12	36	0.05	0.07	0.13	2.6	3	15	1	20	0	3	9	1	6	0	2.4	7.0
run53	12	14	0.09	0.14	0.09	2.1	1	5	6	4	0	0	4	5	1	0	1.6	4.9
run54	12	22	0.09	0.14	0.11	2.3	1	14	12	13	0	0	8	10	9	0	2.1	5.5
run55	12	29	0.09	0.13	0.12	2.5	5	22	12	14	0	2	15	7	7	0	2.0	6.9
run56	12	37	0.09	0.13	0.13	2.6	3	21	6	33	1	2	12	4	14	0	3.2	8.7
run57	12	15	0.18	0.26	0.07	2.3	0	7	9	11	0	0	5	5	3	0	1.3	6.1
run58	12	23	0.18	0.26	0.10	2.4	0	32	19	23	1	0	18	14	7	1	2.4	7.4
run59	12	31	0.18	0.25	0.11	2.6	0	29	32	51	1	0	19	19	13	0	2.1	10.4
run60	12	38	0.18	0.24	0.12	2.7	5	30	21	67	1	3	21	14	24	0	2.3	9.5
run61	12	16	0.27	0.36	0.07	2.4	1	9	20	26	0	1	7	14	15	0	2.0	5.5
run62	12	24	0.27	0.36	0.09	2.6	0	34	41	36	3	0	22	24	18	0	1.9	7.3
run63	12	32	0.27	0.35	0.11	2.6	1	42	41	73	2	0	25	24	24	1	2.4	11.8
run64	12	40	0.27	0.33	0.11	2.8	1	41	49	77	2	0	24	34	19	0	1.8	7.5
run65	12	14	0.05	0.08	0.16	1.8	3	6	4	3	0	2	2	4	1	0	2.6	4.0
run66	12	21	0.05	0.07	0.15	2.1	1	7	0	6	0	1	7	0	2	0	2.2	6.3
run67	12	29	0.05	0.07	0.17	2.2	3	8	1	8	0	1	6	1	6	0	2.6	6.8
run68	12	36	0.05	0.07	0.21	2.2	6	20	5	13	0	1	13	3	4	0	2.4	5.8
run69	12	14	0.09	0.15	0.15	1.8	1	10	8	6	1	0	6	5	4	1	1.6	6.6
run70	12	22	0.09	0.14	0.15	2.1	3	17	12	13	0	2	7	9	6	0	1.2	6.9
run71	12	29	0.09	0.13	0.16	2.3	1	15	7	17	0	1	12	4	8	0	2.1	10.5
run72	12	37	0.09	0.13	0.19	2.3	4	33	10	24	3	1	20	6	13	1	3.1	7.7
run73	12	15	0.18	0.27	0.11	2.0	0	11	15	11	0	0	5	11	6	0	1.5	8.2

TABLE 1
MODEL PROPERTIES AND DYNAMICAL PROPERTIES OF BSSs.

run74	12	23	0.18	0.26	0.14	2.2	3	16	22	19	1	0	11	16	7	0	2.6	7.2
run75	12	30	0.18	0.24	0.14	2.4	1	29	21	42	1	1	19	17	14	1	2.1	6.7
run76	12	38	0.18	0.25	0.17	2.4	3	36	37	52	2	3	21	28	18	1	1.9	10.0
run77	12	16	0.28	0.37	0.09	2.2	1	23	24	19	0	0	14	15	9	0	2.0	8.2
run78	12	24	0.27	0.36	0.12	2.3	0	21	37	41	1	0	12	24	20	1	2.2	7.1
run79	12	32	0.27	0.35	0.13	2.5	2	27	46	73	5	2	20	31	26	0	2.1	8.4
run80	12	40	0.27	0.34	0.17	2.5	2	48	55	59	4	0	23	37	19	1	1.8	7.1
run81	12	14	0.05	0.08	0.37	1.4	1	10	5	1	0	0	7	2	0	0	1.5	2.9
run82	12	21	0.05	0.08	0.36	1.6	1	10	2	4	0	1	5	2	0	0	1.5	5.1
run83	12	28	0.05	0.08	0.42	1.7	8	21	2	1	0	4	16	1	1	0	2.1	5.2
run84	12	36	0.05	0.07	0.42	1.8	4	20	3	7	1	1	16	3	3	0	1.6	6.3
run85	12	14	0.09	0.16	0.33	1.4	0	8	6	0	0	0	7	4	0	0	1.4	4.8
run86	12	22	0.09	0.15	0.34	1.6	1	21	18	6	0	1	13	15	3	0	1.2	10.5
run87	12	29	0.09	0.14	0.36	1.7	4	22	13	11	1	1	17	6	4	0	1.8	5.6
run88	12	36	0.09	0.14	0.40	1.8	10	31	22	12	3	6	23	17	3	0	1.8	6.5
run89	12	15	0.18	0.28	0.21	1.7	0	27	13	5	0	0	19	7	0	0	1.6	5.2
run90	12	23	0.18	0.27	0.25	1.8	2	31	34	16	2	0	23	23	8	0	1.4	7.5
run91	12	30	0.18	0.26	0.30	1.9	1	25	31	26	1	0	18	19	9	0	1.5	6.7
run92	12	38	0.18	0.25	0.30	2.0	5	39	60	38	5	1	24	40	15	1	1.7	7.8
run93	12	16	0.27	0.38	0.15	1.9	0	15	32	9	0	0	10	23	2	0	1.6	6.5
run94	12	24	0.27	0.37	0.21	1.9	1	26	49	19	1	0	15	29	8	1	1.6	5.8
run95	12	32	0.27	0.36	0.24	2.0	2	36	64	43	3	1	20	49	13	2	1.3	7.5
run96	12	40	0.27	0.35	0.27	2.1	2	57	82	53	5	1	31	62	22	4	1.8	7.5
run97	11	14	0.05	0.10	1.26	0.9	2	17	10	1	1	1	12	7	1	0	1.3	3.8
run98	11	21	0.05	0.10	1.71	1.0	3	12	8	0	0	0	8	7	0	0	1.6	3.0
run99	11	28	0.05	0.09	2.04	1.0	4	18	9	1	1	2	11	8	0	1	0.9	4.0
run100	11	36	0.04	0.08	2.23	1.1	9	49	5	3	0	4	40	4	0	0	1.3	6.4
run101	11	14	0.09	0.18	1.29	0.9	0	15	15	6	0	0	9	8	2	0	1.0	4.6
run102	11	22	0.09	0.17	1.33	1.1	3	33	21	3	0	2	21	17	2	0	1.4	6.1
run103	12	36	0.09	0.16	1.88	1.1	5	52	22	5	0	5	37	12	0	0	1.2	5.0
run104	12	15	0.18	0.31	0.77	1.1	3	17	37	6	3	1	10	30	2	1	1.3	6.6
run105	12	23	0.18	0.30	1.13	1.1	1	38	62	7	1	1	27	46	3	0	1.2	6.6
run106	12	30	0.18	0.29	1.07	1.3	2	61	74	8	2	1	43	62	2	2	1.2	7.1
run107	12	38	0.18	0.27	0.93	1.4	3	70	66	7	5	2	43	52	5	0	1.3	7.5
run108	12	15	0.27	0.41	0.46	1.3	0	21	67	6	1	0	12	42	1	0	1.2	5.5
run109	12	23	0.27	0.40	0.67	1.3	0	44	99	3	5	0	30	67	1	1	1.3	6.2
run110	12	32	0.27	0.39	0.65	1.5	1	70	128	10	5	1	42	85	3	2	1.5	6.7
run111	12	40	0.27	0.36	0.65	1.6	5	60	112	14	4	2	38	71	4	3	1.3	7.9
run112	10	14	0.05	0.10	2.02	0.8	1	11	6	3	2	0	8	3	1	2	0.4	6.1
run113	10	21	0.05	0.10	4.29	0.7	2	23	12	0	0	1	19	8	0	0	0.9	3.6
run114	11	14	0.09	0.18	1.37	0.9	2	24	17	0	1	0	12	9	0	0	0.6	5.0
run115	10	22	0.09	0.18	4.07	0.8	5	34	33	0	4	1	23	17	0	1	0.9	4.3
run116	11	14	0.18	0.31	0.93	1.0	1	27	45	4	3	0	19	34	0	1	1.0	3.8
run117	10	23	0.18	0.32	2.33	0.9	4	40	70	1	1	2	25	49	0	1	1.1	6.0
run118	9	38	0.18	0.28	2.33	1.1	4	73	117	2	7	2	50	82	0	3	1.1	5.7
run119	11	15	0.27	0.42	0.84	1.1	2	30	76	7	0	0	17	47	2	0	0.9	4.3
run120	11	23	0.27	0.41	1.25	1.1	1	42	112	1	6	0	24	79	0	3	1.0	5.5
run121	10	32	0.27	0.40	1.62	1.1	1	53	108	4	6	0	39	78	2	2	1.1	5.9

NOTE. — t_{cl} denotes the cluster age in Gyr, M denotes the total cluster mass at t_{cl} in $10^4 M_{\odot}$, f_b denotes the overall binary fraction in the cluster, $f_{b,c}$ denotes the binary fraction in the core, ρ_c denotes the core density in $10^4 M_{\odot} \text{pc}^{-3}$, and r_c denotes the core radius in pc. N_{BSS} and $N_{BSS,c}$ denote the numbers of BSSs in the whole cluster, and within r_c , respectively. The time since formation for the BSSs is denoted by t_{age} in Gyr. COLL, MTB, and SE denote BSSs formed via stellar collisions, stable mass-transfer in a binary, and binary stellar evolution driven mergers, respectively. SS, BS, and BB denote collisions via single-single, binary-single, and binary-binary encounters, respectively. Med, and Max denote the median t_{age} and the maximum t_{age} for each model.